

Evolutionary dynamics of taxonomic structure

Michael Foote

Biol. Lett. 2012 **8**, doi: 10.1098/rsbl.2012.0780 first published online 31 August 2012

Subject collections

Articles on similar topics can be found in the following collections

[evolution](#) (598 articles)

[palaeontology](#) (71 articles)

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

Correction

Biol. Lett. **8**, 135–138 (23 February 2012; Published online 24 August 2011) (doi:10.1098/rsbl.2011.0521)

Evolutionary dynamics of taxonomic structure

Michael Foote

There was a misprint in [table 2](#) of this paper; the corrected table is given here. All calculations in the original paper used the correct figures, so the conclusions of the paper are not affected.

Table 2. Summary statistics on molluscan species and genera from the Cenozoic of New Zealand. Rate estimates ± 1 s.e. based on bootstrap resampling of stratigraphic ranges.

| higher taxon | NF _s | NL _s | NF _g | γ^* | μ^* |
|--------------|-----------------|-----------------|-----------------|------------------|------------------|
| Bivalvia | 589 | 434 | 206 | 0.54 ± 0.017 | 1.13 ± 0.028 |
| Gastropoda | 1502 | 1213 | 415 | 0.38 ± 0.010 | 1.12 ± 0.014 |